

Precipitation of water from aqueous mixtures with addition of hydrophilic ions

Ryuichi Okamoto and Akira Onuki

Department of Physics, Kyoto University, Kyoto 606-8502, Japan

(Dated: July 11, 2011)

We examine phase separation in aqueous mixtures at fixed amounts of hydrophilic monovalent ions. When water is the minority component, preferential solvation can stabilize water domains enriched with ions. This ion-induced precipitation occurs in wide ranges of the temperature and the average composition where the solvent would be in one-phase states without ions. The volume fraction of such water domains is decreased to zero as the interaction parameter χ (dependent on the temperature) is decreased toward a critical value for each average composition.

PACS numbers: 82.45.Gj, 61.20.Qg, 64.75.Cd, 81.16.Dn

In fluid mixtures containing water, phase separation behavior can be drastically changed by a small amount of a salt [1]. More strikingly, many groups have observed long-lived heterogeneities (sometimes extending over a few micrometers) in one-phase states [2] and a third phase visible as a thin plate at a liquid-liquid interface in two-phase states [3]. Very recently, mesophases with lamellar or onion structures have been found for an antagonistic salt composed of hydrophilic and hydrophobic ions [4]. Dramatic ion effects are ubiquitous in various soft matters. For example, in polyelectrolytes, the phase behavior sensitively depend on the degree of ionization and the composition for mixture solvents [5] and large-scale heterogeneities have been observed [6]. In these phenomena, the solvation interaction among ions and polar molecules should play a major role together with the Coulomb interaction among charges [7]. Recently some theoretical efforts have been made to elucidate the solvation effects in phase transitions in electrolytes and polyelectrolytes [8–10]. In this Letter, we consider hydrophilic monovalent ion pairs such as Na^+ and Cl^- in a binary mixture of water and a less polar component (oil or alcohol) and examine ion-induced precipitation.

Neglecting the electrostatic interaction but accounting for the solvation interaction, we first consider a binary mixture in a cell with a fixed volume V . Here ions constitute the third component with density $n(\mathbf{r})$. The volume fractions of water, oil, and ions are written as $\phi(\mathbf{r})$ and $\phi'(\mathbf{r})$, and $v_I n(\mathbf{r})$, respectively, where v_I is the ionic volume. If the two solvent species have the same molecular volume v_0 , their densities are ϕ/v_0 and ϕ'/v_0 . The space-filling condition reads $\phi + \phi' + v_I n = 1$. The ion volume fraction is assumed to be small or $v_I n \ll 1$, which is easily satisfied for small ions with $v_I \ll v_0$. In this Letter we thus set $\phi' = 1 - \phi$, which simplifies the calculations.

When the ions have a strong preference of water over oil, we set up the free energy as

$$\frac{F}{T} = \int d\mathbf{r} \left[\frac{f(\phi)}{T} + \frac{C}{2} |\nabla \phi|^2 + n \ln(nv_0) - gn\phi \right], \quad (1)$$

where C is a positive constant and g is a positive parameter representing the ion preference of water. The space integral is within the cell. The Boltzmann constant will

be set equal to unity and then the temperature T represents the thermal energy of a particle. The free energy density $f(\phi)$ is taken to be the simple form,

$$v_0 f/T = \phi \ln \phi + (1 - \phi) \ln(1 - \phi) + \chi \phi(1 - \phi), \quad (2)$$

where χ is the interaction parameter dependent on T and its mean-field critical value is 2 without ions. We fix the total particle numbers of the three components as

$$\bar{n} = \int d\mathbf{r} n/V, \quad \bar{\phi} = \int d\mathbf{r} \phi/V, \quad (3)$$

In equilibrium the homogeneity of the ion chemical potential $\delta F/\delta n$ yields

$$n = \bar{n} e^{g\phi} / \langle e^{g\phi} \rangle, \quad (4)$$

where $\langle e^{g\phi} \rangle = \int d\mathbf{r} e^{g\phi}/V$ is the space average of $e^{g\phi}$. Substitution of Eq.(4) into F in Eq.(1) gives

$$\frac{F}{T} = \int d\mathbf{r} \left[\frac{f(\phi)}{T} + \frac{C}{2} |\nabla \phi|^2 \right] + V \bar{n} \ln[\bar{n} v_0 / \langle e^{g\phi} \rangle]. \quad (5)$$

In equilibrium the chemical potential difference $h = \delta F/\delta \phi$ for the composition is also homogeneous. Here,

$$h = f'(\phi) - TC \nabla^2 \phi - Tgn \quad (6)$$

where $f' = \partial f/\partial \phi$. Around a planar interface varying along the z axis, we obtain $TC(d\phi/dz)^2 = 2H(\phi)$, where

$$H(\phi) = f(\phi) - f(\phi_\alpha) - T(n - n_\alpha) - h(\phi - \phi_\alpha). \quad (7)$$

We suppose coexistence of a water-rich phase α and an oil-rich phase β with $\Delta\phi = \phi_\alpha - \phi_\beta > 0$. The compositions and the ion densities in the two phases are written as ϕ_α , ϕ_β , n_α , and n_β , respectively. The volume fraction of the phase α is denoted by γ_α . Since that of the phase β is $\gamma_\beta = 1 - \gamma_\alpha$, we have

$$\gamma_\alpha = (\bar{\phi} - \phi_\beta)/\Delta\phi = (\bar{n} - n_\beta)/\Delta n, \quad (8)$$

where $\Delta n = n_\alpha - n_\beta$. From Eq.(4) the ratio of the bulk ion densities is written as $n_\alpha/n_\beta = \exp(g\Delta\phi) \gg 1$ from

Eq.(4), where we assume $g\Delta\phi \gg 1$. Neglecting the surface free energy, we express F as

$$\frac{F}{V} = \gamma_\alpha f_\alpha + \gamma_\beta f_\beta - T\bar{n} \log[(\gamma_\alpha e^{g\phi_\alpha} + \gamma_\beta e^{g\phi_\beta})/\bar{n}v_0], \quad (9)$$

where $f_\alpha = f(\phi_\alpha)$ and $f_\beta = f(\phi_\beta)$. We minimize this F with respect to γ_α , ϕ_α , and ϕ_β at fixed $\bar{\phi}$ to obtain

$$h = f'_\alpha - Tg n_\alpha = f'_\beta - Tg n_\beta, \quad (10)$$

$$f_\alpha - f_\beta - T\Delta n = h\Delta\phi, \quad (11)$$

where $f'_\alpha = f'(\phi_\alpha)$ and $f'_\beta = f'(\phi_\beta)$. These equations also follow from Eqs.(6) and (7).




FIG. 1: Compositions ϕ_α and ϕ_β (left) and γ_α , $\gamma_\alpha n_\alpha/\bar{n}$, and n_α/\bar{n} (right) vs χ , where $\bar{n} = 6 \times 10^{-4} v_0^{-1}$ and $g = 11$. For $\bar{\phi} = 0.35$ (top), ϕ_α continuously changes for $\chi > \chi_p = 0.687$. For $\bar{\phi} = 0.46$ (bottom), ϕ_α jumps at $\chi \cong 2$, where $\chi_p = 1.834$. Shown also are the coexistence curve without ions and the spinodal curve with ions. The latter follows from a shift of the spinodal curve without ions by $g^2 v_0 \bar{n}/2$ to the left [8].

Our main finding is that there appears a precipitation branch of $\phi = \phi_\alpha \cong 1$ in the region $\chi < 2$ for $g \gg 1$. In Eq.(10) we set $f'_\alpha \cong v_0^{-1} T[-\log(1 - \phi_\alpha) - \chi]$ to obtain

$$1 - \phi_\alpha \cong A_\alpha \exp[-gG(\phi_\beta)], \quad (12)$$

where $A_\alpha = \exp(\chi + v_0 f'_\beta/T)$. Here we introduce

$$G(\phi) = -v_0[f(\phi) + (1 - \phi)f'(\phi)]/T \\ = -\log \phi - \chi(1 - \phi)^2, \quad (13)$$

where the second line follows from Eq.(2). By setting $f_\alpha \cong 0$ in Eq.(11) we also find $v_0 n_\alpha = G(\phi_\beta)$. Outside the spinodal curve without ions, we notice $dG/d\phi =$




FIG. 2: $\chi = \chi_p(\bar{\phi})$ for $v_0 \bar{n} = 4 \times 10^{-4}$, 2×10^{-4} , and 5×10^{-5} . Precipitation occurs for $\bar{\phi} < \phi_p$ (arrows). Shown also are the coexistence and spinodal curves without ions.

$-v_0(1 - \phi)\partial^2 f/\partial\phi^2 < 0$, which leads to $G(\phi) > 0$ from $G(1) = 0$. Thus the right hand side of Eq.(12) is small or $\phi_\alpha \cong 1$ for large g . The equation for ϕ_β is given by

$$v_0 n_\alpha = G(\phi_\beta) = v_0 \bar{n} \Delta\phi / [\bar{\phi} - \phi_\beta + e^{-g\Delta\phi} \Delta\phi] \quad (14)$$

where we may set $\Delta\phi \cong 1 - \phi_\beta$. As $\gamma_\alpha = (\bar{\phi} - \phi_\beta)/\Delta\phi$ decreases to zero, ϕ_β and n_α increase up to $\bar{\phi}$ and $v_0 \bar{n} e^{g(1 - \bar{\phi})}$, respectively. Remarkably, ϕ_α and ϕ_β depend on $\bar{\phi}$. From the second line of Eq.(13), the precipitation branch exists only for $\chi > \chi_p(\bar{\phi})$. The lower bound is

$$\chi_p = [-\log(1 - \bar{\phi}) - v_0 \bar{n} e^{g(1 - \bar{\phi})}]/(1 - \bar{\phi})^2, \quad (15)$$

where $\gamma_\alpha \rightarrow 0$ as $\chi \rightarrow \chi_p$.

To easily understand the mathematics, let us focus on the case $\gamma_\alpha \ll 1$, where $f_\alpha \cong 0$ and $f_\beta \cong f(\bar{\phi}) - f'_\beta(1 - \bar{\phi})\gamma_\alpha$ in Eq.(9). Treating γ_α as an order parameter, we write the deviation $\Delta F = F(\gamma_\alpha) - F(0)$ of the free energy from the one-phase value $F(0)$ as

$$\Delta F/TV \cong A_1 \gamma_\alpha - \bar{n} \log(1 + B_1 \gamma_\alpha), \quad (16)$$

where $A_1 = v_0^{-1} G(\bar{\phi}) + \bar{n} g(1 - \bar{\phi})$ and $B_1 = e^{g\Delta\phi} - 1 \cong e^{g\Delta\phi}$. For $w \equiv \bar{n} B_1/A_1 > 1$, ΔF has a negative minimum given by $-TV A_1(w \log w - w + 1)/B_1 < 0$ attained at

$$\gamma_\alpha = (w - 1)/B_1 \cong v_0 \bar{n}/G(\bar{\phi}) - e^{-g(1 - \bar{\phi})}, \quad (17)$$

which is consistent with Eq.(14) in the limit $\phi_\beta \rightarrow \bar{\phi}$. The condition $w > 1$ is equivalent to $\chi > \chi_p$ for $g \gg 1$.

Figure 1 gives the phase diagrams in the ϕ - χ plane with $\bar{n} = 6 \times 10^{-4} v_0^{-1}$ and $g = 11$. In the first case of $\bar{\phi} = 0.35$, ϕ_α changes continuously and is minimum at $\chi = 2.05$, where $\chi_p = 0.687$ and the maximum of n_α is $0.381 v_0^{-1}$ at $\chi = \chi_p$. In the second case of $\bar{\phi} = 0.46$, where $\chi_p = 1.834$ and $n_\alpha = 0.0842 v_0^{-1}$ at $\chi = \chi_p$, we find that ϕ_α changes discontinuously along a hysteresis loop in the range $1.953 < \chi < 1.967$. In equilibrium, F is minimized and the resultant discontinuous transition

is at $\chi = 1.957$. In Fig.2, we display curves of $\chi = \chi_p(\bar{\phi})$ for three values of \bar{n} with $g = 10$. Each curve assumes a minimum at small $\bar{\phi}$ far away from the coexistence curve without ions. With increasing $\bar{\phi}$, the precipitation branch shrinks and disappears as $\bar{\phi} \rightarrow \phi_p$, where ϕ_p is a critical composition less than 0.5. As functions of $\bar{\phi}$ and χ , we show γ_α in Fig.3 in the continuous case of $g = 10$ and ϕ_α in Fig.4 in the discontinuous case of $g = 11$.

Without the electrostatic interaction, the surface tension of our system is expressed as the integral $\sigma = \int dz CT(d\phi/dz)^2$ around an interface varying along the z axis [8]. Use of Eq.(7) gives

$$\sigma = (2CT)^{1/2} \int_{\phi_\beta}^{\phi_\alpha} d\phi H(\phi)^{1/2}, \quad (18)$$

where C is assumed to be a constant. In Fig.5, we display the function $[2H(\phi)v_0/T]^{1/2}$ for $g = 11$. Here $H(\phi) \cong f''(\phi_\beta)(\phi - \phi_\beta)^2/2$ as $\phi \rightarrow \phi_\beta$ with $f'' = d^2f/d\phi^2$. Thus we obtain $\sigma \sim TC(\Delta\phi)^2/2\xi$, where $\xi = (f''/CT)^{-1/2}$ is the correlation length at $\phi = \phi_\beta \cong \bar{\phi}$.




FIG. 3: Volume fraction of the water-rich phase γ_α as a function of χ and $\bar{\phi}$. for $\bar{n} = 2 \times 10^{-4} v_0^{-1}$ and $g = 10$. In this case γ_α decreases continuously to zero as $\chi \rightarrow \chi_p$.

Including the electrostatic interaction, we next consider aqueous mixtures containing a hydrophilic monovalent salt. The cation and anion densities are written as n_1 and n_2 , whose total amounts are fixed as

$$\int d\mathbf{r} n_1 = \int d\mathbf{r} n_2 = V\bar{n}/2. \quad (19)$$

The electric potential Φ satisfies the Poisson equation $\nabla \cdot \varepsilon \nabla \Phi = -4\pi e(n_1 - n_2)$, where the dielectric constant $\varepsilon(\phi)$ can depend on ϕ . The free energy F reads [8]

$$\begin{aligned} \frac{F}{T} = & \int d\mathbf{r} \left[\frac{f(\phi)}{T} + \frac{C}{2} |\nabla \phi|^2 + \frac{\varepsilon |\nabla \Phi|^2}{8\pi T} \right. \\ & \left. + \sum_{i=1,2} [n_i \ln(n_i v_0) - g_i n_i \phi] \right]. \end{aligned} \quad (20)$$

The ion chemical potentials due to solvation, written as $\mu_{\text{sol}}^i(\phi)$, strongly depend on ϕ . The interaction terms




FIG. 4: (Color on line) Composition of the water-rich phase ϕ_α for $v_0 \bar{n} = 6 \times 10^{-4}$ and $g = 11$. In this case a discontinuous transition occurs for $\chi \cong 1.95$ and $\bar{\phi} > 0.395$ (in red). One-phase states are realized for $\chi < \chi_p$ or for $\bar{\phi} > \phi_p = 0.473$ and $\chi < 2$ (in white). For $\chi > 2$, ϕ_α is little affected by ions.




FIG. 5: $[2H(\phi)v_0/T]^{1/2}$ vs ϕ , vanishing at $\phi = \phi_\alpha$ and ϕ_β , where $\bar{\phi} = 0.46$, $\bar{n} = 6 \times 10^{-4} v_0^{-1}$, and $g = 11$.

($\propto g_i$) in F follow for the linear forms $\mu_{\text{sol}}^i(\phi) = \mu_0^i - T g_i \phi$ (where the first terms are irrelevant constants). This linear dependence is adopted to gain the physical consequences in the simplest manner. For each ion species i , the solvation-chemical-potential difference between the two phases is given by $\Delta\mu_{\alpha\beta}^i = T g_i \Delta\phi$, which is the Gibbs transfer free energy in electrochemistry [11]. In aqueous solutions, $g_i \gg 1$ for hydrophilic small ions, while $g_i < 0$ for hydrophobic ions [8, 11]. We minimize F with respect to n_i under Eq.(19) to obtain

$$n_i = n_i^0 \exp[g_i \phi \mp e\Phi/T], \quad (21)$$

where $-$ is for $i = 1$, $+$ is for $i = 2$, and $n_i^0 = \bar{n}/[2 \int d\mathbf{r} \exp(g_i \phi \mp e\Phi/T)]$. The composition profile is determined by the homogeneity of $\delta F/\delta \phi$. Here we neglect the image interaction, whose role is reduced compared to that of the solvation interaction for not small ion densities [8, 12].

For $g_1 = g_2 = g$, we have $n_1 = n_2 = n/2$ and $\Phi = 0$,




FIG. 6: Left: Numerical droplet radius d/a vs χ (bold line) together with the theoretical curve without the surface free energy (dotted line). Right: Normalized potential $e\Phi(r)/T$ and water volume fraction $\phi(r)$ (top), and normalized ion densities $v_0n_1(r)$ and $v_0n_2(r)$ (bottom), where $\chi = 1.7$. Here $g_1 = 15$, $g_2 = 7$, $\bar{\phi} = 0.35$, and $v_0\bar{n} = 6 \times 10^{-4}$.

so F in Eq.(20) reduces to F in Eq.(1). For $g_1 \neq g_2$, an electric double layer appears at the interface with a potential difference $\Delta\Phi = T(g_1 - g_2)\Delta\phi/2e$ across it, but the bulk phase relations (10) and (11) still hold with

$$g = (g_1 + g_2)/2. \quad (22)$$

In Fig.6, we give numerical results for $g_1 = 15$, $g_2 = 7$, $\bar{\phi} = 0.35$, and $v_0\bar{n} = 6 \times 10^{-4}$. The space unit is $a = v_0^{1/3}$. We suppose a spherical water-rich droplet with radius d placed at the center of a spherical cell with radius $R = 600a$. Then $\gamma_\alpha = (d/R)^3$. The dielectric constant is of the form $\epsilon = 40(1 + \phi)$. We also set $aC = \chi$ and $e^2/T = 120a$. In the left, the droplet disappears at $d = 26.0a$, where w in Eq.(17) is 1.08. This critical radius follows if the bulk free energy ΔF in Eq.(9) is equated with the minims of the surface free energy $4\pi\sigma d^2$. In the right, we

set $\chi = 1.7$ to display the profiles of $\Phi(r)$, $\phi(r)$, $n_1(r)$, and $n_2(r)$. We obtain $\phi_\alpha = 0.993$ and $n_\alpha = 0.352v_0^{-1}$ within the droplet and $\phi_\beta = 0.349$ and $n_\beta = 2.55 \times 10^{-4}v_0^{-1}$ outside it. In Fig.6, the potential $\Phi(r)$ relaxes to the one-dimensional solution of the nonlinear Poisson-Boltzmann equation (dotted line) with $\Delta\Phi = 3T/e$ [8]. Here n_α and n_β are the bulk values of $n = n_1 + n_2$.

In future we should explain the experimental findings of large-scale heterogeneities [2]. We note that one-phase states are metastable outside the spinodal curve with ions ($\chi < 1/2\bar{\phi}(1 - \bar{\phi}) - g^2v_0\bar{n}/2$) in Fig.1. Thus precipitation from a one-phase state should be triggered by some impurities and/or hydrophilic walls. We also note that the wetting transition of aqueous mixtures is much influenced by the ion-induced precipitation mechanism.

Experiments are informative, where the temperature, the water volume fraction, and the salt amount are varied. We mention an experiment by Leunissen *et al.* [13], where micron-sized water droplets containing ions formed a crystal in an oil with low dielectric constant ($\epsilon_{\text{oil}} = 4 - 10$) without a surfactant. Graaf *et al.* [14] ascribed its origin to the screened Coulomb interaction among droplets. We also propose experiments of the salting-out effect of polyelectrolytes in water-alcohol [5], where the degree of ionization much increases with accumulation of water around the polymers [10].

Acknowledgments

This work was supported by Grant-in-Aid for Scientific Research on Priority Area ‘‘Soft Matter Physics’’ from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

-
- [1] E.L. Eckfeldt and W.W. Lucasse, J. Phys. Chem. **47**, 164 (1943); B.J. Hales *et al.*, J. Phys. Chem. **70**, 3970 (1966); V. Balevicius and H. Fuess, Phys. Chem. Chem. Phys. **1**, 1507 (1999); M. Misawa *et al.*, J. Phys. and Chem. of Solids, **60**, 1301 (1999).
 - [2] G. W. Euliss and C. M. Sorensen, J. Chem. Phys. **80**, 4767 (1984); A. F. Kostko *et al.*, Phys. Rev. E **70**, 026118 (2004); M. Wagner *et al.*, Phys. Chem. Chem. Phys. **6**, 580 (2004); C. Yang, W. Li, and C. Wu, J. Phys. Chem. B **108**, 11866 (2004); M. Sedlak, J. Phys. Chem. B **110**, 4329, 4339, 13976 (2006).
 - [3] J. Jacob *et al.*, Phys. Chem. Chem. Phys. **3**, 829 (2001).
 - [4] K. Sadakane *et al.*, J. Phys. Soc. Jpn., **76**, 113602 (2007); Phys. Rev. Lett., **103**, 167803 (2009).
 - [5] P. G. Arscott *et al.*, Biopolymers, **36**, 345 (1995); A. Hultgren and D. C. Rau, Biochemistry **43**, 8272 (2004); C. Stanley and D. C. Rau, Biophys. J. **91**, 912 (2006).
 - [6] For example, see B. D. Ermi and E. J. Amis, Macromolecules **31**, 7378 (1998).
 - [7] J. N. Israelachvili, *Intermolecular and Surface Forces* (Academic Press, London, 1991).
 - [8] A. Onuki, Phys. Rev. E **73**, 021506 (2006); J. Chem. Phys. **128**, 224704 (2008).
 - [9] T. Araki and A. Onuki, J. Phys.: Condens. Matter **21**, 424116 (2009).
 - [10] A. Onuki and R. Okamoto, J. Phys. Chem. B, **113**, 3988 (2009); R. Okamoto and A. Onuki, J. Chem. Phys. **131**, 094905 (2009).
 - [11] L. Q. Hung, J. Electroanal. Chem. **115**, 159 (1980); T. Osakai and K. Ebina, J. Phys. Chem. B **102**, 5691 (1998). For water-nitrobenzene at $T \cong 300\text{K}$, the Gibbs transfer free energy was estimated as 17 for Na^+ and as 19 for Cl^- per ion in units of $k_B T$, so $g_i \sim 20$ for this case.
 - [12] Y. Levin and J. E. Flores-Mena, Europhys. Lett. **56**, 187 (2001).
 - [13] M. E. Leunissen *et al.*, Proc. Natl. Acad. Sci. U.S.A. **104**, 2585 (2007); Phys. Chem. Chem. Phys. **9**, 6405 (2007).
 - [14] J. de Graaf *et al.*, J. Chem. Phys. **129**, 194701 (2008).